


**IN THE CLAIMS:**

1. – 17. (Cancelled):

18. (New): A method of treating inflammatory conditions comprising administering to a mammal in need thereof an effective amount of a cyclic compound selected from the group consisting of cyclopentane, cyclohexane, cycloheptane, monosaccharide, disaccharide, trisaccharide, tetrasaccharide, piperidine, tetrahydrothiopyran, 5-oxotetrahydrothiopyran, 5,5-dioxotetrahydrothiopyran, tetrahydroselenopyran, tetrahydrofuran, pyrrolidine, tetrahydrothiophene, 5-oxotetrahydrothiophene, 5,5-dioxotetrahydrothiophene, tetrahydroselenophene, benzene, cumene, mesitylene, naphthalene and phenanthrene, in which said cyclic compound is substituted by at least three vicinal phosphorus containing radicals of the formula:



wherein

$\text{V}^1$  to  $\text{V}^4$  are independently  $\text{Y}_{m6}^8 \text{T}_{o3} \text{U}$ ;

$\text{T}_{o1}$  to  $\text{T}_{o3}$  are independently  $(\text{CH}_2)_n$ ,  $\text{CH}=\text{CH}$ , or  $\text{CH}_2\text{CH}=\text{CHCH}_2$ ;

$o1$  to  $o3$  are independently 0 or 1;

n is 0 to 4;

U is  $R^1Y^9$ ,  $CY^{10}Y^{11}R^2$ ,  $SY^{12}Y^{13}Y^{14}R^3$ ,  $PY^{15}Y^{16}Y^{17}R^4R^5$ ,

$Y^{18}PY^{19}Y^{20}Y^{21}R^6R^7$ ,  $CH_2NO_2$ ,  $NHSO_2R^8$ , or  $NHCY^{22}Y^{23}R^9$ ;

m1 to m7 are independently 0 or 1;

$Y^1$  to  $Y^{23}$  are independently  $NR^{10}$ ,  $NOR^{11}$ , O, or S;

and where  $R^1$  to  $R^{11}$  are independently

i) hydrogen;

ii) a straight or branched saturated or unsaturated alkyl group containing 1-22 carbon atoms;

iii) a saturated, unsaturated aromatic or non-aromatic homo- or heterocyclic group containing 3-22 carbon atoms and 0-5 heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur;

iv) a straight or branched saturated or unsaturated alkyl group containing 1-22 carbon atoms substituted with a saturated or unsaturated aromatic or non-aromatic homo- or heterocyclic group containing 3-22 carbon and 0-5 heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur;


v) an aromatic or non-aromatic homo-or heterocyclic group containing 3-22 carbon and 0-5 heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur which aromatic or non-aromatic homo-or heterocyclic group is substituted with a straight or branched saturated or unsaturated group containing 1-22 carbon atoms;

whereby

said groups in ii-v are unsubstituted or are substituted by 1-6 of the following groups: hydroxy, alkoxy, aryloxy, acyloxy, carboxy, alkoxy carbonyl, alkoxy carbonyloxy, aryloxy carbonyl, aryloxy carbonyloxy, carbamoyl, fluoro, chloro, bromo, azido, cyano, oxo, oxa, amino, imino, alkylamino, arylamino, acylamino, arylazo, nitro, alkylthio or alkylsulfonyl.

19. (New): The method according to Claim 18 wherein the inflammatory conditions is rheumatoid arthritis.

20. (New): The method according to Claim 18 or 19 wherein the phosphorus-containing radicals have the following formula:



wherein

$\text{V}^1$  and  $\text{V}^2$  are OH,  $(\text{CH}_2)_p\text{OH}$ , COOH,  $\text{CONH}_2$ , CONOH,  $(\text{CH}_2)_p\text{COOH}$ ,  $(\text{CH}_2)_p\text{CONH}_2$ ,  $(\text{CH}_2)_p\text{CONOH}$ ,  $(\text{CH}_2)_p\text{SO}_3\text{H}$ ,  $(\text{CH}_2)_p\text{SO}_3\text{NH}_2$ ,  $(\text{CH}_2)_p\text{NO}_2$ ,  $(\text{CH}_2)_p\text{PO}_3\text{H}_2$ ,  $\text{O}(\text{CH}_2)_p\text{OH}$ ,  $\text{O}(\text{CH}_2)_p\text{COOH}$ ,  $\text{O}(\text{CH}_2)_p\text{CONH}_2$ ,  $\text{O}(\text{CH}_2)_p\text{CONOH}$ ,  $\text{O}(\text{CH}_2)_p\text{SO}_3\text{H}$ ,  $\text{O}(\text{CH}_2)_p\text{SO}_3\text{NH}_2$ ,  $\text{O}(\text{CH}_2)_p\text{NO}_2$ ,  $\text{O}(\text{CH}_2)_p\text{PO}_3\text{H}_2$  or  $\text{CF}_2\text{COOH}$ ; and


$p$  is 1 to 4.

21. (New): The method according to Claim 18 or 19 wherein the phosphorus-containing radicals are phosphate groups.

22. (New): The method according to Claim 18 or 19 wherein the cyclic compound is a monosaccharide.

23. (New): The method according to Claim 22 wherein the monosaccharide is D/L-ribose, D/L-arabinose, D/L-xylose, D/L-lyxose, D/L-allose, D/L-altrose, D/L-glucose, D/L-mannose, D/L-gulose, D/L-idose, D/L-galactose, D/L-talose, D/L-ribulose, D/L-xylulose, D/L-psicose, D/L-sorbose, D/L-tagatose, or D/L-fructose.

24. (New): The method according to Claim 22 wherein the monosaccharide is substituted with three phosphorus-containing radicals having the following formula:



wherein

$\text{V}^1$  and  $\text{V}^2$  are OH,  $(\text{CH}_2)_p\text{OH}$ , COOH, CONH<sub>2</sub>, CONOH,  $(\text{CH}_2)_p\text{COOH}$ ,  $(\text{CH}_2)_p\text{CONH}_2$ ,  $(\text{CH}_2)_p\text{CONOH}$ ,  $(\text{CH}_2)_p\text{SO}_3\text{H}$ ,  $(\text{CH}_2)_p\text{SO}_3\text{NH}_2$ ,  $(\text{CH}_2)_p\text{NO}_2$ ,  $(\text{CH}_2)_p\text{PO}_3\text{H}_2$ ,  $\text{O}(\text{CH}_2)_p\text{OH}$ ,  $\text{O}(\text{CH}_2)_p\text{COOH}$ ,  $\text{O}(\text{CH}_2)_p\text{CONH}_2$ ,  $\text{O}(\text{CH}_2)_p\text{CONOH}$ ,  $\text{O}(\text{CH}_2)_p\text{SO}_3\text{H}$ ,  $\text{O}(\text{CH}_2)_p\text{SO}_3\text{NH}_2$ ,  $\text{O}(\text{CH}_2)_p\text{NO}_2$ ,  $\text{O}(\text{CH}_2)_p\text{PO}_3\text{H}_2$  or  $\text{CF}_2\text{COOH}$ ; and  
 $p$  is 1 to 4.


25. (New): The method according to Claim 24 wherein the phosphorous containing radicals are phosphate groups.

26. (New): The method according to Claim 18 or 19 wherein the cyclic compound administered to the mammal is selected from the group consisting of mannose-2,3,4-trisphosphate, rhamnose-2,3,4-trisphosphate, galactose- 2,3,4-trisphosphate, methyl-6-O-butyl- $\alpha$ -D-mannopyranoside-2,3,4-trisphosphate, 1,5-anhydro-D-arabinitol-2,3,4-trisphosphate, fructose-2,3,4-trisphosphate, 1,2-O-ethylene- $\beta$ -D-fructopyranoside-2,3,4-trisphosphate, cyclohexane-1,2,3-triol trisphosphate, 1,5-dideoxy-1,5-iminoarabinitol-2,3,4-trisphosphate, altrose-2,3,4-trisphosphate, or methyl-6-O-butyl- $\alpha$ -D-altropyranoside 2,3,4-trisphosphate.

27. (New): The method according to Claim 18 or 19 wherein the compound is administered by parenteral or non-parenteral administration.

28. (New): The method according to Claim 18 or 19 wherein the effective amount ranges from about 0.1 to about 100 mg per kg body weight of the mammal.

29. (New): A method of treating tissue repair conditions comprising administering to a mammal in need thereof an effective amount of a cyclic compound selected from the group consisting of cyclopentane, cyclohexane, cycloheptane, inositol, monosaccharide, disaccharide, trisaccharide, tetrasaccharide, piperidine, tetrahydrothiopyran, 5-oxotetrahydrothiopyran, 5,5-dioxotetrahydrothiopyran, tetrahydroselenopyran, tetrahydrofuran, pyrrolidine, tetrahydrothiophene, 5-oxotetrahydrothiophene, 5,5-dioxotetrahydrothiophene, tetrahydroselenophene, benzene, cumene, mesitylene, naphthalene and phenanthrene, in which said cyclic compound is substituted by at least three vicinal phosphorus containing radicals of the formula:



wherein

$\text{V}^1$  to  $\text{V}^4$  are independently  $\text{Y}^8_{\text{m}6}\text{T}_{03}\text{U}$ ;

$T_{01}$  to  $T_{03}$  are independently  $(CH_2)_n$ ,  $CH=CH$ , or  $CH_2CH=CHCH_2$ ;

$o_1$  to  $o_3$  are independently 0 or 1;

$n$  is 0 to 4;

$U$  is  $R^1Y^9m_7$ ,  $CY^{10}Y^{11}R^2$ ,  $SY^{12}Y^{13}Y^{14}R^3$ ,  $PY^{15}Y^{16}Y^{17}R^4R^5$ ,

$Y^{18}PY^{19}Y^{20}Y^{21}R^6R^7$ ,  $CH_2NO_2$ ,  $NHSO_2R^8$ , or  $NHCY^{22}Y^{23}R^9$ ;

$m_1$  to  $m_7$  are independently 0 or 1;

$Y^1$  to  $Y^{23}$  are independently  $NR^{10}$ ,  $NOR^{11}$ , O, or S;

and where  $R^1$  to  $R^{11}$  are independently

i) hydrogen;

ii) a straight or branched saturated or unsaturated alkyl group containing 1-22 carbon atoms;

iii) a saturated, unsaturated aromatic or non-aromatic homo- or heterocyclic group containing 3-22 carbon atoms and 0-5 heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur;

iv) a straight or branched saturated or unsaturated alkyl group containing 1-22 carbon atoms substituted with a saturated or unsaturated aromatic or non-aromatic homo- or heterocyclic group containing 3-22 carbon and 0-5 heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur;

v) an aromatic or non-aromatic homo-or heterocyclic group containing 3-22 carbon and 0-5 heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur which aromatic or non-aromatic homo-or heterocyclic group is substituted with a straight or branched saturated or unsaturated group containing 1-22 carbon atoms;


whereby

said groups in ii-v are unsubstituted or are substituted by 1-6 of the following groups: hydroxy, alkoxy, aryloxy, acyloxy, carboxy, alkoxycarbonyl, alkoxycarbonyloxy, aryloxy carbonyl, aryloxy carbonyloxy, carbamoyl, fluoro, chloro, bromo, azido, cyano, oxo, oxa, amino, imino, alkylamino, arylamino, acylamino, arylazo, nitro, alkylthio or alkylsulfonyl.



30. (New): A method according to Claim 29 wherein the tissue repair condition is wound healing, matrix formation, collagen synthesis or scar formation.

31. (New): The method according to Claim 29 or 30 wherein the phosphorus-containing radicals have the following formula:



wherein

$\text{V}^1$  and  $\text{V}^2$  are OH,  $(\text{CH}_2)_p\text{OH}$ , COOH, CONH<sub>2</sub>, CONOH,  $(\text{CH}_2)_p\text{COOH}$ ,  $(\text{CH}_2)_p\text{CONH}_2$ ,  $(\text{CH}_2)_p\text{CONOH}$ ,  $(\text{CH}_2)_p\text{SO}_3\text{H}$ ,  $(\text{CH}_2)_p\text{SO}_3\text{NH}_2$ ,  $(\text{CH}_2)_p\text{NO}_2$ ,  $(\text{CH}_2)_p\text{PO}_3\text{H}_2$ ,  $\text{O}(\text{CH}_2)_p\text{OH}$ ,  $\text{O}(\text{CH}_2)_p\text{COOH}$ ,  $\text{O}(\text{CH}_2)_p\text{CONH}_2$ ,  $\text{O}(\text{CH}_2)_p\text{CONOH}$ ,  $\text{O}(\text{CH}_2)_p\text{SO}_3\text{H}$ ,  $\text{O}(\text{CH}_2)_p\text{SO}_3\text{NH}_2$ ,  $\text{O}(\text{CH}_2)_p\text{NO}_2$ ,  $\text{O}(\text{CH}_2)_p\text{PO}_3\text{H}_2$  or  $\text{CF}_2\text{COOH}$ ; and

$p$  is 1 to 4.


32. (New): The method according to Claim 29 or 30 wherein the phosphorus-containing radicals are phosphate groups.

33. (New): The method according to Claim 29 or 30 wherein the cyclic compound is a monosaccharide.

34. (New): The method according to Claim 33 wherein the monosaccharide is D/L-ribose, D/L-arabinose, D/L-xylose, D/L-lyxose, D/L-allose, D/L-altrose, D/L-glucose, D/L-mannose,

D/L-gulose, D/L-idose, D/L-galactose, D/L-talose, D/L-ribulose, D/L-xylulose, D/L-psicose, D/L-sorbose, D/L-tagatose, or D/L-fructose.

35. (New): The method according to Claim 33 wherein the monosaccharide is substituted with three phosphorus-containing radicals having the following formula:



wherein


$\text{V}^1$  and  $\text{V}^2$  are OH,  $(\text{CH}_2)_p\text{OH}$ , COOH, CONH<sub>2</sub>, CONOH,  $(\text{CH}_2)_p\text{COOH}$ ,  $(\text{CH}_2)_p\text{CONH}_2$ ,  $(\text{CH}_2)_p\text{CONOH}$ ,  $(\text{CH}_2)_p\text{SO}_3\text{H}$ ,  $(\text{CH}_2)_p\text{SO}_3\text{NH}_2$ ,  $(\text{CH}_2)_p\text{NO}_2$ ,  $(\text{CH}_2)_p\text{PO}_3\text{H}_2$ ,  $\text{O}(\text{CH}_2)_p\text{OH}$ ,  $\text{O}(\text{CH}_2)_p\text{COOH}$ ,  $\text{O}(\text{CH}_2)_p\text{CONH}_2$ ,  $\text{O}(\text{CH}_2)_p\text{CONOH}$ ,  $\text{O}(\text{CH}_2)_p\text{SO}_3\text{H}$ ,  $\text{O}(\text{CH}_2)_p\text{SO}_3\text{NH}_2$ ,  $\text{O}(\text{CH}_2)_p\text{NO}_2$ ,  $\text{O}(\text{CH}_2)_p\text{PO}_3\text{H}_2$  or  $\text{CF}_2\text{COOH}$ ; and  $p$  is 1 to 4.

36. (New): The method according to Claim 35 wherein the phosphorous containing radicals is a phosphate group.

37. (New): The method according to Claim 29 or 30 wherein the cyclic compound is inositol.

38. (New): The method according to Claim 37 wherein the inositol is alloinositol, cisinositol, ipiinositol, D/L-chiroinositol, scylloinositol, myoinositol, mycoinositol or neoinositol.

39. (New): The method according to Claim 38 wherein the inositol is substituted with three phosphorus-containing radicals having the following formula:



wherein

$\text{V}^1$  and  $\text{V}^2$  are OH,  $(\text{CH}_2)_p\text{OH}$ , COOH, CONH<sub>2</sub>, CONOH,  $(\text{CH}_2)_p\text{COOH}$ ,  $(\text{CH}_2)_p\text{CONH}_2$ ,  $(\text{CH}_2)_p\text{CONOH}$ ,  $(\text{CH}_2)_p\text{SO}_3\text{H}$ ,  $(\text{CH}_2)_p\text{SO}_3\text{NH}_2$ ,  $(\text{CH}_2)_p\text{NO}_2$ ,  $(\text{CH}_2)_p\text{PO}_3\text{H}_2$ ,  $\text{O}(\text{CH}_2)_p\text{OH}$ ,  $\text{O}(\text{CH}_2)_p\text{COOH}$ ,  $\text{O}(\text{CH}_2)_p\text{CONH}_2$ ,  $\text{O}(\text{CH}_2)_p\text{CONOH}$ ,  $\text{O}(\text{CH}_2)_p\text{SO}_3\text{H}$ ,  $\text{O}(\text{CH}_2)_p\text{SO}_3\text{NH}_2$ ,  $\text{O}(\text{CH}_2)_p\text{NO}_2$ ,  $\text{O}(\text{CH}_2)_p\text{PO}_3\text{H}_2$  or  $\text{CF}_2\text{COOH}$ ; and p is 1 to 4.

40. (New): The method according to Claim 35 wherein the phosphorous containing radicals are phosphate groups.


41. (New): The method according to Claim 29 or 30 wherein the cyclic compound administered to the mammal is selected from the group consisting of myoinositol-1,2,6-trisphosphate, mannose-2,3,4-trisphosphate, rhamnose-2,3,4-trisphosphate, galactose-2,3,4-trisphosphate, methyl-6-O-butyl- $\alpha$ -D-mannopyranoside-2,3,4-trisphosphate, 1,5-anhydro-D-arabinitol-2,3,4-trisphosphate, fructose-2,3,4-trisphosphate, 1,2-O-ethylene- $\beta$ -D-fructopyranoside-2,3,4-trisphosphate, cyclohexane-1,2,3-triol trisphosphate, 1,5-dideoxy-1,5-

iminoarabinitol-2,3,4-trisphosphate, altrose-2,3,4-trisphosphate, or methyl-6-O-butyl- $\alpha$ -D-altropyranoside 2,3,4-trisphosphate.

42. (New): The method according to Claim 29 or 30 wherein the compound is administered by parenteral or non-parenteral administration.

43. (New): The method according to Claim 29 or 20 wherein the effective amount ranges from about 0.1 to about 100 mg per kg body weight of the mammal.

44. (New): A method of treating infectious conditions comprising administering to a mammal in need thereof an effective amount of a cyclic compound selected from the group consisting of cyclopentane, cyclohexane, cycloheptane, inositol, monosaccharide, disaccharide, trisaccharide, tetrasaccharide, piperidine, tetrahydrothiopyran, 5-oxotetrahydrothiopyran, 5,5-dioxotetrahydrothiopyran, tetrahydroselenopyran, tetrahydrofuran, pyrrolidine, tetrahydrothiophene, 5-oxotetrahydrothiophene, 5,5-dioxotetrahydrothiophene, tetrahydroselenophene, benzene, cumene, mesitylene, naphthalene and phenanthrene, in which said cyclic compound is substituted by at least three vicinal phosphorus containing radicals of the formula:



wherein

$V^1$  to  $V^4$  are independently  $Y^8_{m6}T_{o3}U$ ;

$T_{o1}$  to  $T_{o3}$  are independently  $(CH_2)_n$ ,  $CH=CH$ , or  $CH_2CH=CHCH_2$ ;

$o1$  to  $o3$  are independently 0 or 1;

$n$  is 0 to 4;

$U$  is  $R^1Y^9_{m7}$ ,  $CY^{10}Y^{11}R^2$ ,  $SY^{12}Y^{13}Y^{14}R^3$ ,  $PY^{15}Y^{16}Y^{17}R^4R^5$ ,

$Y^{18}PY^{19}Y^{20}Y^{21}R^6R^7$ ,  $CH_2NO_2$ ,  $NHSO_2R^8$ , or  $NHCY^{22}Y^{23}R^9$ ;

$m1$  to  $m7$  are independently 0 or 1;

$Y^1$  to  $Y^{23}$  are independently  $NR^{10}$ ,  $NOR^{11}$ ,  $O$ , or  $S$ ;

and where  $R^1$  to  $R^{11}$  are independently

i) hydrogen;

ii) a straight or branched saturated or unsaturated alkyl group containing 1-22 carbon atoms;

iii) a saturated, unsaturated aromatic or non-aromatic homo- or heterocyclic group containing 3-22 carbon atoms and 0-5 heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur;

iv) a straight or branched saturated or unsaturated alkyl group containing 1-22 carbon atoms substituted with a saturated or unsaturated aromatic or non-aromatic homo- or heterocyclic group containing 3-22 carbon and 0-5 heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur;


v) an aromatic or non-aromatic homo-or heterocyclic group containing 3-22 carbon and 0-5 heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur which aromatic or non-aromatic homo-or heterocyclic group is substituted with a straight or branched saturated or unsaturated group containing 1-22 carbon atoms;

whereby

said groups in ii-v are unsubstituted or are substituted by 1-6 of the following groups: hydroxy, alkoxy, aryloxy, acyloxy, carboxy, alkoxycarbonyl, alkoxycarbonyloxy, aryloxy carbonyl, aryloxy carbonyloxy, carbamoyl, fluoro, chloro, bromo, azido, cyano, oxo, oxa, amino, imino, alkylamino, arylamino, acylamino, arylazo, nitro, alkylthio or alkylsulfonyl.

45. (New): A method according to Claim 44 wherein the infectious condition is trypanosomiasis.

46. (New): The method according to Claim 44 or 45 wherein the phosphorus-containing radicals have the following formula:



wherein


$\text{V}^1$  and  $\text{V}^2$  are OH,  $(\text{CH}_2)_p\text{OH}$ , COOH, CONH<sub>2</sub>, CONOH,  $(\text{CH}_2)_p\text{COOH}$ ,  $(\text{CH}_2)_p\text{CONH}_2$ ,  $(\text{CH}_2)_p\text{CONOH}$ ,  $(\text{CH}_2)_p\text{SO}_3\text{H}$ ,  $(\text{CH}_2)_p\text{SO}_3\text{NH}_2$ ,  $(\text{CH}_2)_p\text{NO}_2$ ,  $(\text{CH}_2)_p\text{PO}_3\text{H}_2$ ,  $\text{O}(\text{CH}_2)_p\text{OH}$ ,  $\text{O}(\text{CH}_2)_p\text{COOH}$ ,  $\text{O}(\text{CH}_2)_p\text{CONH}_2$ ,  $\text{O}(\text{CH}_2)_p\text{CONOH}$ ,  $\text{O}(\text{CH}_2)_p\text{SO}_3\text{H}$ ,  $\text{O}(\text{CH}_2)_p\text{SO}_3\text{NH}_2$ ,  $\text{O}(\text{CH}_2)_p\text{NO}_2$ ,  $\text{O}(\text{CH}_2)_p\text{PO}_3\text{H}_2$  or  $\text{CF}_2\text{COOH}$ ; and  
 $p$  is 1 to 4.

47. (New): The method according to Claim 44 or 45 wherein the phosphorus-containing radicals are phosphate groups.

48. (New): The method according to Claim 44 or 45 wherein the cyclic compound is a monosaccharide.

49. (New): The method according to Claim 48 wherein the monosaccharide is D/L-ribose, D/L-arabinose, D/L-xylose, D/L-lyxose, D/L-allose, D/L-altrose, D/L-glucose, D/L-mannose, D/L-gulose, D/L-idose, D/L-galactose, D/L-talose, D/L-ribulose, D/L-xylulose, D/L-psicose, D/L-sorbose, D/L-tagatose, or D/L-fructose.

50. (New): The method according to Claim 48 wherein the monosaccharide is substituted with three phosphorus-containing radicals having the following formula:



wherein

$\text{V}^1$  and  $\text{V}^2$  are OH,  $(\text{CH}_2)_p\text{OH}$ , COOH, CONH<sub>2</sub>, CONOH,  $(\text{CH}_2)_p\text{COOH}$ ,  $(\text{CH}_2)_p\text{CONH}_2$ ,  $(\text{CH}_2)_p\text{CONOH}$ ,  $(\text{CH}_2)_p\text{SO}_3\text{H}$ ,  $(\text{CH}_2)_p\text{SO}_3\text{NH}_2$ ,  $(\text{CH}_2)_p\text{NO}_2$ ,  $(\text{CH}_2)_p\text{PO}_3\text{H}_2$ ,  $\text{O}(\text{CH}_2)_p\text{OH}$ ,  $\text{O}(\text{CH}_2)_p\text{COOH}$ ,  $\text{O}(\text{CH}_2)_p\text{CONH}_2$ ,  $\text{O}(\text{CH}_2)_p\text{CONOH}$ ,  $\text{O}(\text{CH}_2)_p\text{SO}_3\text{H}$ ,  $\text{O}(\text{CH}_2)_p\text{SO}_3\text{NH}_2$ ,  $\text{O}(\text{CH}_2)_p\text{NO}_2$ ,  $\text{O}(\text{CH}_2)_p\text{PO}_3\text{H}_2$  or  $\text{CF}_2\text{COOH}$ ; and p is 1 to 4.


51. (New): The method according to Claim 50 wherein the phosphorous containing radicals are phosphate groups.

52. (New): The method according to Claims 44 or 45 wherein the cyclic compound is inositol.



53. (New): The method according to Claim 52 wherein the inositol is alloinositol, cisinisitol, ipiinositol, D/L-chiroinositol, scylloinositol, myoinositol, mycoinositol or neoinositol.

54. (New): The method according to Claim 53 wherein the inositol is substituted with three phosphorus-containing radicals having the following formula:



wherein

$\text{V}^1$  and  $\text{V}^2$  are OH,  $(\text{CH}_2)_p\text{OH}$ , COOH, CONH<sub>2</sub>, CONOH,  $(\text{CH}_2)_p\text{COOH}$ ,  $(\text{CH}_2)_p\text{CONH}_2$ ,  $(\text{CH}_2)_p\text{CONOH}$ ,  $(\text{CH}_2)_p\text{SO}_3\text{H}$ ,  $(\text{CH}_2)_p\text{SO}_3\text{NH}_2$ ,  $(\text{CH}_2)_p\text{NO}_2$ ,  $(\text{CH}_2)_p\text{PO}_3\text{H}_2$ ,  $\text{O}(\text{CH}_2)_p\text{OH}$ ,  $\text{O}(\text{CH}_2)_p\text{COOH}$ ,  $\text{O}(\text{CH}_2)_p\text{CONH}_2$ ,  $\text{O}(\text{CH}_2)_p\text{CONOH}$ ,  $\text{O}(\text{CH}_2)_p\text{SO}_3\text{H}$ ,  $\text{O}(\text{CH}_2)_p\text{SO}_3\text{NH}_2$ ,  $\text{O}(\text{CH}_2)_p\text{NO}_2$ ,  $\text{O}(\text{CH}_2)_p\text{PO}_3\text{H}_2$  or  $\text{CF}_2\text{COOH}$ ; and  
 $p$  is 1 to 4.

55. (New): The method according to Claim 54 wherein the phosphorous containing radicals is a phosphate group.

56. (New): The method according to Claim 44 or 45 wherein the cyclic compound administered to the mammal is selected from the group consisting of myoinositol-1,2,6-trisphosphate, mannose-2,3,4-trisphosphate, rhamnose-2,3,4-trisphosphate, galactose- 2,3,4-

triphosphate, methyl-6-O-butyl- $\alpha$ -D-mannopyranoside-2,3,4-triphosphate, 1,5-anhydro-D-arabinitol-2,3,4-triphosphate, fructose-2,3,4-triphosphate, 1,2-O-ethylene- $\beta$ -D-fructopyranoside-2,3,4-triphosphate, cyclohexane-1,2,3-triol triphosphate, 1,5-dideoxy-1,5-iminoarabinitol-2,3,4-triphosphate, altrose-2,3,4-triphosphate, or methyl-6-O-butyl- $\alpha$ -D-altropyranoside 2,3,4-triphosphate.

57. (New): The method according to Claim 44 or 45 wherein the compound is administered by parenteral or non-parenteral administration.

58. (New): The method according to Claim 44 or 45 wherein the growth factor modulating effective amount ranges from about 0.1 to about 100 mg per kg body weight of the mammal.